

Working Paper Series

COVID-19: US Federal accountability for entry, spread, and inequities

W. P. Hanage¹, C. Testa², J. T. Chen², L. Davis³, E. Pechter³, M. Santillana⁴ and N. Krieger²

October 5, 2020

HCPDS Working Paper Volume 20, Number 2

The views expressed in this paper are those of the author(s) and do not necessarily reflect those of the Harvard Center for Population and Development Studies.

Affiliations

- 1. Center for Communicable Disease Dynamics, Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA
- 2. Department of Social and Behavioral Sciences, Harvard T.H. Chan School of Public Health, Boston, MA
- 3. Consultant, Boston, MA
- 4. Computational Health Informatics Program, Boston Children's Hospital, Boston, MA

Corresponding Author

Nancy Krieger, PhD Professor of Social Epidemiology American Cancer Society Clinical Research Professor Department of Social and Behavioral Sciences Harvard T.H. Chan School of Public Health 677 Huntington Avenue Boston, MA 02115 Email: <u>nkrieger@hsph.harvard.edu</u>

Abstract

The United States (US) has been among those nations most severely affected by the first—and subsequent—phases of the pandemic of COVID-19 disease caused by SARS-CoV-2. With only 4% of the worldwide population, the US has seen about 22% of COVID-19 deaths. Despite formidable advantages in resources and expertise, presently the per capita mortality rate is over 585/million, respectively 2.4 and 5 times higher compared to Canada and Germany. As we enter Fall 2020, the US is enduring ongoing outbreaks across large regions of the country. Moreover, within the US, an early and persistent feature of the pandemic has been the disproportionate impact on populations already made vulnerable by racism and dangerous jobs, inadequate wages, and unaffordable housing, and this is true for both the headline public health threat and the additional disastrous economic impacts. In this article we assess the impact of missteps by the Federal Government in three specific areas: the introduction of the virus to the US and the establishment of community transmission; the lack of national COVID-19 workplace standards and lack of personal protective equipment (PPE) for workplaces as represented by complaints to the Occupational Safety and Health Administration (OSHA) which we find are correlated with deaths 17 days later (=0.845); and the total excess deaths in 2020 to date, which already total more than 230,000 and exhibit severe inequities in race/ethnicity including among younger age groups.

COVID-19: US Federal accountability for entry, spread, and inequities

Authors:

W. P. Hanage¹, C. Testa², J. T. Chen², L. Davis³, E. Pechter³, M. Santillana⁴ and N. Krieger²

¹Center for Communicable Disease Dynamics, Department of Epidemiology, Harvard T. H. Chan School of Public Health, Boston, MA.

²Department of Social and Behavioral Sciences, Harvard T.H. Chan School of Public Health, Boston, MA.

³Consultant, Boston, MA.

⁴Computational Health Informatics Program, Boston Children's Hospital, Boston, MA

Abstract

The United States (US) has been among those nations most severely affected by the first—and subsequent—phases of the pandemic of COVID-19 disease caused by SARS-CoV-2. With only 4% of the worldwide population, the US has seen about 22% of COVID-19 deaths. Despite formidable advantages in resources and expertise, presently the per capita mortality rate is over 585/million, respectively 2.4 and 5 times higher compared to Canada and Germany. As we enter Fall 2020, the US is enduring ongoing outbreaks across large regions of the country. Moreover, within the US, an early and persistent feature of the pandemic has been the disproportionate impact on populations already made vulnerable by racism and dangerous jobs, inadequate wages, and unaffordable housing, and this is true for both the headline public health threat and the additional disastrous economic impacts. In this article we assess the impact of missteps by the Federal Government in three specific areas: the introduction of the virus to the US and the establishment of community transmission; the lack of national COVID-19 workplace standards and lack of personal protective equipment (PPE) for workplaces as represented by complaints to the Occupational Safety and Health Administration (OSHA) which we find are correlated with deaths 17 days later (ρ =0.845); and the total excess deaths in 2020 to date, which already total more than 230,000 and exhibit severe inequities in race/ethnicity including among younger age groups.



Living graffiti at Jamaica Pond (Boston, MA): early May, while COVID-19 lockdown underway – "We cannot direct the wind, but we can adjust the sail" (photograph taken by N. Krieger) The United States (US) has been among those nations most severely affected by the first—and subsequent—phases of the pandemic of COVID-19 disease caused by SARS-CoV-2. With only 4% of the worldwide population, the US has seen about 22% of COVID-19 deaths. Despite formidable advantages in resources and expertise, presently the per capita mortality rate is over 585/million, respectively 2.4 and 5 times higher compared to Canada and Germany [1, 2]. As we enter Fall 2020, the US is enduring ongoing outbreaks across large regions of the country. Moreover, within the US, an early and persistent feature of the pandemic has been the disproportionate impact on populations already made vulnerable by racism and dangerous jobs, inadequate wages, and unaffordable housing, and this is true for both the headline public health threat and the additional disastrous economic impacts [3-8].

The newly issued federal *Healthy People 2030* framework (released on August 18, 2020) [9], sets the nation's objectives for the next decade, with the first two overarching goals being: (1) "Attain healthy, thriving lives and well-being free of preventable disease, injury, and premature death" and (2) "Eliminate health disparities, achieve health equity, and attain health literacy to improve the health and well-being of all." The third is: "Create social, physical and economic environments that promote attaining the full potential for health and well-being for all" [10]. In the case of COVID-19, the US has missed the mark on all three goals.

In the following we document critical federal missteps in protecting the nation, with an eye towards guiding a better response. We start with how community transmission in the US became established in the spring.

Control of borders: entry of the virus

To prevent or delay establishment of the virus, many countries have implemented symptom screening at ports of entry, sometimes together with quarantine of travelers. Such measures have enabled Singapore, Hong Kong, and Taiwan to avoid large early outbreaks, despite having been at high risk of introductions due to the volume of travelers from Wuhan (Hubei province, China), the epicenter of the pandemic. In the US, control of borders, including in relation to public health, rests with the federal government, an authority assumed since the founding of the country [11-15]. The administration implemented a travel ban affecting only non-US travelers from China on January 31, 2020, without any mandatory symptom screening at entry or quarantine of travelers, despite the virus at this time already known to be present in Italy, Iran, Spain, Germany, Finland, and the United Kingdom. Similarly, selective restrictions on travel from Europe were only instituted a full 6 weeks later, on March 11, 2020, at which point 830 deaths had been reported in Italy alone (by contrast, on January 31, 2020 China had reported 259 deaths). None of these restrictions applied to returning US citizens or permanent residents, despite their potential exposure to SARS-CoV-2 [16].

Hence there were ongoing opportunities for introductions of the virus to the US from Europe. Consistent with this, phylogenetic studies have found repeatedly that the great majority of SARS-CoV-2 introductions to the US were viral lineages circulating in Europe [62]. A study of the genomic epidemiology of the early pandemic in the Boston area estimated a total of 25 independent importations from Europe, 18 of which occurred prior to March 28 [17]. Imports included a superspreading event, a conference, on February 26-27 leading to more than 90 known infections among attendees or their contacts. The index cases were two international business travelers. This introduction is estimated to be the direct ancestor of 35% of sampled genomes not associated with the conference, including samples from outbreaks in homeless shelters, so it is reasonable to suggest that this introduction alone hastened the establishment of the virus in the region and the scale of the subsequent surge of infections. As has been previously shown, by early March the most likely source of introduction to a region was from elsewhere in the United States [18]. This shows that any action to delay or prevent introductions would need to be at the national level, and while the window for effective action was brief, it was missed.

Symptom screening alone is unlikely to provide a significant barrier to a virus like SARS-CoV-2 that is capable of presymptomatic transmission [63], but may be an effective component of mitigation if supplemented by a coherent central strategy to delay case importation. While like-for-like comparisons of government responses to COVID-19 should be interpreted with caution, the benefits of such an approach are well-illustrated by Australia: like the US, Australia is a large and well-developed economy with major metropolitan cities, substantial international travel, and wide variations in population density. Australia had reported 300 confirmed cases on March 15 when it imposed a closure of its borders to non-Australian citizens, which was augmented by a mandatory 14-day, supervised quarantine in a hotel for all international arrivals (including Australian citizens), aggressive testing and contact tracing, and population-wide adherence to social distancing guidelines [19]. Following these local containment measures Australia saw no explosive COVID-19 case surges through the Southern hemisphere winter. Despite a recent burst of domestic transmission in the state of Victoria that resulted in the immediate re-imposition of a statewide lockdown and interstate movement restrictions [20], Australia presently reports 25.6 deaths/million persons, in comparison with the US report of 585 deaths/million persons [21]. This is the difference between establishing effective national border policies and failing to do so.

Lack of occupational health protections and community spread

The pandemic has presented different risks and rates of exposure to different sectors of US society, at different times and in different places. Health care and other essential workers who continued to work while much of the country was shut down, or who work in close quarters or have contact with the public as part of their job—e.g., in transportation, critical manufacturing, food and agriculture, grocery stores, and pharmacies—have faced a greater risk of exposure and infection from COVID-19. [22-26]. Despite the designation of these workers as "essential," an important federal failure has been the lack of standards requiring systematic collection of data on COVID-19 cases or deaths by industry and occupation. Nevertheless, information available from case reports, state health departments, media reports, and other sources show: large numbers of cases among health care workers and first responders; major outbreaks in meatpacking, poultry, and agriculture plants; outbreaks among prison staff (above and beyond inmates); and numerous cases in transportation, warehousing, and other essential industries, as well as in public-facing occupations (e.g., grocery stores and retail) [27]. People infected at work expose their household members—and if they are low-wage workers, they are more likely to live in crowded housing with inadequate ventilation, which increases the risk of transmission and decreases the options for people who are ill to self-isolate [7-8, 28-31].

The federal government could have limited transmission of SARS-CoV-2 at the workplace in multiple ways. One is through its unparalleled purchasing power and ability to invoke the Defense

Production Act and ensure equitable supply and distribution of PPE. However supplying PPE was delegated to a variety of actors: state and city governments, large hospital chains, and in some cases small networks of clinics [13]. Second, the federal government could have established mandatory universal paid sick leave for those unable to work due to Covid-19, but it did not. The congressional mandate for 80 hours of paid COVID-related leave covered only some groups of workers. Third, the federal government could have mandated standards for occupational exposures, but it failed to act, even as some US states have done so. To date, the federal Occupational Safety and Health Administration (OSHA) has not issued any emergency or permanent standard specific to COVID-19 exposure at the workplace; additionally, as of August 13, 2020, OSHA, which oversees enforcement of OSHA standards in 38 states, had issued only four citations related to COVID-19 [32]. Moreover, the total number of federal OSHA inspections (of any kind) during 2020 has been reduced by two-thirds, compared to the same period in prior years [33].

Several of the 22 states with approved state OSHA plans have taken stronger regulatory and enforcement action than federal OSHA to protect workers from exposure to SARS-CoV-2. For example, Virginia issued a comprehensive emergency temporary standard on SARS-CoV-2 on July 27, 2020, and the state of Michigan has issued multiple executive orders on worker protection from COVID-19 enforced through the Michigan State OSHA plan [34-35]. The California, Nevada, and Minnesota state OSHA plans have all issued numerous citations under other existing standards to protect workers from COVID-19. However, in most states little regulatory or enforcement action has been taken by either the federal government or the state to address workplace exposures to the virus [36]. While there have been few federal OSHA citations, worker complaints to OSHA that raise concerns about workplace conditions and exposure to COVID-19 can serve to estimate hazards of exposures and risks of infection as reported by workers themselves. To our knowledge, these data have not been analyzed in relation to the population burdens of COVID-19. Figure 1 shows the volume of OSHA complaints (federal and state combined), broken down by four industries (manufacturing, retail, healthcare and social assistance, and other). The total number of national complaints is shown, as are the numbers of complaints for the four main geographical regions of the country. Mortality data from confirmed cases is also shown. Two findings stand out:

(1) The curve of the OSHA complaint data resembles an epidemic curve, and (2) the complaints predate COVID-19 deaths. Pearson correlation coefficients indicate that nationwide for the period Feb 1–Aug 22 there is a correlation between OSHA complaints and COVID-19 mortality with a lag of 17 days (ρ = 0.845). The variation in rates by geographic region and industry is substantial and mirrors the course of the pandemic. Regional correlations between complaints and reported cases and deaths (Fig 1) suggest complaints reflect local responses to awareness of pandemic activity, and that the nationwide correlation is not driven by any one region. These patterns point to a lack of learning from experiences across states, along with the lack of enforceable federal standards.

We cannot determine the extent to which patterns of OSHA complaints reflect growing awareness of the pandemic and the need for adequate protection, or whether they are more direct indicators of failures that causally contributed to workplace, family, and community spread. But findings do suggest that expressed worker concerns may be an indicator of real risks, and failure to respond is a missed opportunity to intervene to mitigate disease transmission in the workplace and, in turn, the community at large. While there is not a consistent definition of "essential workers," data suggest that people in this category disproportionately are low-income, are people of color, rely on public transportation, and live in crowded housing [7-8, 23-25]. They also have more underlying disabilities and less access to care underscores the importance of mitigating exposures to SARs-CoV-2 in the workplace. Work is a key exposure risk, and to date, federal policies regarding COVID-19 and workplaces have chiefly aimed to limit the liability of employers, rather than to protect employees through, for example, mandating that employers develop an infection control plan[64]. In the following section we consider the consequences in terms of excess all-cause mortality.

The outcome: total deaths, inequities in premature mortality, and increasing cases. Beyond the number of confirmed COVID-19 deaths in the US (more than 190,000 to date), estimates of excess deaths (comparing the same calendar period for 2020 to the average annual deaths for 2015-2019 and controlling for population growth) provide a powerful way to gauge the excess mortality due to COVID-19. Excess mortality may be due directly to the infection or be due to delays in seeking care for non-COVID illnesses or injuries; this measure also takes into account potential reductions in mortality (e.g., due to lower air pollution during the economic lockdown) as well as the seasonality of death rates [37-39].

In Table 1, we show the results of four different approaches, ranging from more to less conservative, that have been used to estimate the excess number of deaths and the excess death rate (per 100,000), in relation to US mortality data as of August 16, 2020 [40-43]. No matter which

method is used, the estimated excess number of deaths, ranging from 231,049 to 263,375, is already greater than the outer margin of what the federal government stated, on April 3, 2020, would be the upper limit of the expected death toll, i.e., 200,000 [44-46].

We additionally estimated an annual excess age-standardized death rate of 83.3 per 100,000 p-y based on the difference in age-standardized mortality rates for 2020 vs 2015-2019, (as opposed to scaling the estimated excess death count by the 2020 population as in Table 1). This estimate is nearly double that of the third leading cause of death in the US, i.e., unintentional injuries, for which the annual age-standardized death rate was 48/100,000 p-y in 2018 (the most recent year available) [47]. The excess deaths since the start of the pandemic are already over half of those attributed to cancer for all of 2018 (annual age-standardized death rate 149.1/100,000 p-y) [47]. It should be obvious that given that the pandemic is not over, and transmission continues in much of the country, these numbers can only increase.

Australia is again instructive, as it exhibits very little excess mortality. The most recent report from the Australian Bureau of Statistics states, "55,047 doctor certified deaths occurred between 1 January 2020 and 26 May 2020 and were registered by 30 June. This compares to a baseline average of 53,361 over the past 5 years" [48]. This translates to an excess of only 1686 deaths in Australia. In contrast, the US, with a population 13 times greater, has 154 times more excess deaths.

We must also consider outcomes other than death. While SARS-CoV-2 is a virus that spreads via the respiratory route, it causes a systemic infection with commensurate potential for diverse long

term sequelae, including important outcomes such as stroke in young, otherwise healthy adults [49], and Multisystem Inflammatory Syndrome in Children (MIS-C), a condition similar to Kawasaki syndrome that is associated with pediatric SARS-CoV-2 infection [50]. If an extremely large number of infections are permitted, even rare chronic outcomes will leave a large burden on healthcare after the pandemic. As of August 27, 2020, the number of confirmed cases in the US equaled 6.04 million—almost on par with the population of Massachusetts (6.9 million) [51]. If 0.5% (30,222) develop persistent debilitating symptoms, this would be similar to the number of new cases of cancer per year in Massachusetts (which reported an average of 31,166 per year for 2012-2016) [52]. If 5% of the 6 million cases were so afflicted, the number of persons (300,000) would exceed the 2020 national estimate for the number of US women diagnosed with breast cancer (279,100) [53]. The proportion of long-term effects of one kind or another is unknown, but it is increasingly recognized as a source of serious concern [54].

Finally, despite the initial federal failure to report COVID-19 data by race/ethnicity [6], a combination of specific studies, state reporting, investigative journalism, and data trackers has revealed that a persistent feature of the pandemic has been the existence of racial/ethnic inequities in cases, hospitalizations, and mortality, especially with regard to increased risk among US Black, Latinx, and American Indian/Alaska Native populations compared to the US white non-Hispanic population [3-5, 7-8, 55-56]. What is less appreciated is that racial/ethnic inequities in COVID-19 mortality rates, especially among younger working-age adults, are increasing over time, especially among the Latinx and American Indian/Alaska Native populations. These trends are evident in Figure 2, which we generated using CDC data for February 1 through August 22, 2020 (with 99.2% of deaths assigned a race/ethnicity) [42, 57]. Specifically, the racial/ethnic inequity in

age-standardized mortality rate ratios, compared to white non-Hispanics, increased over time for the five calendar periods displayed (February 1-May 2, May 3-June 6, June 7-July 4, July 5-August 1, and August 2-22) among working age adults (ages 25-64) who were categorized as Black non-Hispanic (p value for trend <0.0001), Hispanic (p < 0.0001), and American Indian/Alaska Native (p <0.0001).The estimates we report, moreover, are likely to be conservative because classification of COVID-19 deaths in part depends on testing, and inequities in access to testing, especially affecting low-income communities of color, would deflate their reported COVID-19 mortality rates [58-59]. Moreover, the elevated mortality risk on the order of 5- to 10-fold among persons under age 65 shown in Figure 2 is unlikely to be explained by pre-existing co-morbidities, because the documented racial/ethnic mortality rate ratios for major chronic conditions (including cardiovascular disease, diabetes, and cancer) among adults under age 65 are around 2-fold or less [56, 60-61].

In closing, we have documented key points at which the US federal government has failed to take the actions and collect the data needed to protect the US population from COVID-19 and the attendant toll on population health and health inequities. Despite the understandable dismay at the state of the pandemic in the United States, it is not too late to make a difference, and all of these failures can and should be reversed by implementation of apt policies. All cases and deaths cannot be prevented—that clearly is not possible with a novel pandemic—but the evidence suggests that ineffective national policies and responses [13], especially as compared to those of other wealthy nations or compared to the intricate preparation and planning by previous administrations of both parties, have been driving the terrible toll of COVID-19 and its inequities in the US. This country—and its political leaders, who bear responsibility—can and must do better. Acknowledgements:

We would like to thank Ziba Cranmer and Gowtham Asokan, with BU SPARK! (Boston University, Massachusetts), for their advice, early on, regarding accessing and exploring the OSHA complaint data, and also thank Peg Seminario (Director of Occupational Safety and Health for the AFL-CIO from 1990-2019) for her expert contributions to interpreting and contextualizing the OSHA data. We would also like to thank Lawson Ung (HSPH) for advice on the nature of the Australian pandemic response. We would like to thank and acknowledge Dr. Mary T. Bassett (Director, François-Xavier Bagnoud (FXB) Center for Health and Human Rights at Harvard University and FXB Professor of the Practice of Health and Human Rights, Harvard T.H. Chan School of Public Health) for her intellectual leadership for prior work, in which she involved NK and JTC, on the importance of analyzing age-specific racial/ethnic inequities in COVID-19 mortality, and which has influenced the approach taken for this paper. We would also like to thank Pamela Waterman (HSPH) for her influence on the paper's analytic orientation and also for her role in facilitating team meetings and communication. References

[1] Schiffman A. nCoV2019.live website. 2020 (<u>https://ncov2019.live/data</u>).

[2] Worldometer—Coronavirus, "Reported Cases and Deaths by Country, Territory, or

Conveyance." 2020 (https://www.worldometers.info/coronavirus/#countries).

[3] Blow C. The racial time bomb in the Covid-19 Crisis. New York Times 2020 April 1

(https://www.nytimes.com/2020/04/01/opinion/coronavirus-black-people.html).

[4] Kendi IX. What the data show: the pandemic seems to be hitting people of color the hardest.

The Atlantic 2020 April 6 (https://www.theatlantic.com/ideas/archive/2020/04/coronavirus-

exposing-our-racial-divides/609526/].

[5] Bailey Z, Barber S, Robinson W, et al. Racism in the time of COVID-19. Interdisciplinary Association for Population Health Science 2020 April 9 (https://iaphs.org/racism-in-the-time-ofcovid-19/).

[6] Krieger N, Gonsalves G, Bassett MT, et al. The fierce urgency of now: closing glaring gaps in US surveillance data on COVID-19. Health Affairs Blog 2020 April 14

(https://www.healthaffairs.org/do/10.1377/hblog20200414.238084/full/].

[7] Diez Roux A. Population health in the time of COVID-19: confirmations and revelations.
 Milbank Quarterly 2020 August 18 (https://onlinelibrary.wiley.com/doi/full/10.1111/1468-0009.12474).

[8] Krieger N. ENOUGH: COVID-19, structural racism, police brutality, plutocracy, climate change and time for health justice, democratic governance, and an equitable, sustainable future. Am J Public Health 2020 August 20:e1-e4.

(https://ajph.aphapublications.org/doi/abs/10.2105/AJPH.2020.305886) (Epub ahead of print).

[9] US Dept of Health and Human Services (US DHHS). Healthy people 2030: building a healthier future for all. 2020 August 18 (https://health.gov/healthypeople).

[10] US DHHS. Healthy people 2030 framework. 2020 August 18

(https://health.gov/healthypeople/about/healthy-people-2030-framework).

[11] Rosen G. A history of public health (Revised expanded ed.). Baltimore, MD: Johns Hopkins University Press, 2015.

[12] United States Federal Security Agency. History of the United States public health service:

1798–1948. Washington, DC: Federal Security Agency, 1948.

[13] Burris S, de Guia S, Gable L, et al. (Eds.). Assessing legal responses to COVID-19. Boston, MA:Public Health Law Watch, 2020 August 21

(https://www.publichealthlawwatch.org/covid19-policy-playbook).

[14] Institute of Medicine. The future of public health. Washington, DC: The National AcademiesPress, 1988 (https://www.nap.edu/catalog/1091/the-future-of-public-health).

[15] Fee E. Public health and the state: the United States. In: Porter D (ed). The History of PublicHealth and the Modern State, pp. 224-275. Amsterdam: Rodopoi, 1994.

[16] Miller G, Dawsey J, Davis AC. One final viral infusion: Trump's move to block travel from

Europe triggered chaos and a surge of passengers from the outbreak's center. Washington Post

2020 May 23 (https://www.msn.com/en-ca/news/world/one-final-viral-infusion-trumps-move-to-

block-travel-from-europe-triggered-chaos-and-a-surge-of-passengers-from-the-outbreaks-

center/ar-BB14v5kG).

[17] Lemieux J, Siddle KJ, Shaw BM, et al. Phylogenetic analysis of SARS-CoV-2 in the Boston area

highlights the role of recurrent importation and superspreading events. 2020 August 25

(https://www.medrxiv.org/content/10.1101/2020.08.23.20178236v1) preprint.

[18] Fauver JR, Petrone ME, Hodcroft EB, et al. Coast-to-coast spread of SARS-CoV-2 during the early epidemic in the United States. Cell 2020;5:990-996.e5

(https://www.sciencedirect.com/science/article/pii/S0092867420304840).

[19] McAnulty JM, Ward K. Suppressing the epidemic in New South Wales. N Engl J Med

2020;382:e74 (https://www.nejm.org/doi/full/10.1056/NEJMc2011592).

[20] Australian Government Dept of Health. Covid-19 in Victoria.

(https://www.health.gov.au/news/health-alerts/novel-coronavirus-2019-ncov-health-

alert/coronavirus-covid-19-current-situation-and-case-numbers/covid-19-in-victoria).

[21] World Health Organization (WHO). WHO coronavirus disease (COVID-19) dashboard.

(https://covid19.who.int).

[22] CDC. Instructions for completing the Human Infection with 2019 Novel Coronavirus (COVID-

19) case report form. Version 1, May 2020. (https://www.cdc.gov/coronavirus/2019-

ncov/downloads/COVID-19-Persons-Under-Investigation-and-Case-Report-Form-Instructions.pdf).

[23] Rogers TN, Rogers CR, VanSant-Webb E, et al. Racial disparities in COVID-19 mortality among

essential workers in the United States. World Medical & Health Policy 2020 August 5

(doi:10.1002/wmh3.358).

[24] Baker MG. Nonrelocatable occupations at increased risk during pandemics: United States,
2018. Am J Public Health 2020;110(8):1126-1132 (doi:10.2105/AJPH.2020.305738) (Epub ahead of print).

[25] Steinberg J, Kennedy ED, Basler C, et al. COVID-19 outbreak among employees at a meat processing facility—South Dakota, March-April 2020. Morb Mortal Wkly Rep 2020;69(31):1015-1019 (doi:10.15585/mmwr.mm6931a2. PMID: 32759914).

[26] Hayden JC, Parkin R. The challenges of COVID-19 for community pharmacists and opportunities for the future. Irish J Psychol Med. 2020;May 21:1-6.

(https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7276501/).

[27] Houlihan CF, Vora N, Byrne T, et al. Pandemic peak SARS-CoV-2 infection and seroconversion rates in London frontline health-care workers. The Lancet 2020;396(10246):e6-e7.

[28] Madewell ZJ, Yang Y, Longini IM, et al. Household transmission of SARS-CoV-2: a systematic review and meta-analysis of secondary attack rate. 2020 July 31

(https://www.medrxiv.org/content/10.1101/2020.07.29.20164590v1). preprint.

[29] Solis J, Franco-Paredes C, Henao-Martínez AF, et al. Structural vulnerability in the U.S.

revealed in three waves of COVID-19. Am J Trop Med Hyg 2020;103(1):25-27

(doi:10.4269/ajtmh.20-0391) (Epub ahead of print).

[30] Chin T, Kahn R, Li R, et al. US-county level variation in intersecting individual, household, and community characteristics relevant to COVID-19 and planning an equitable response: a cross-

sectional analysis. BMJ Open 2020;10(9):e039886

(https://bmjopen.bmj.com/content/10/9/e039886).

[31] Chen JT, Krieger N. Revealing the unequal burden of COVID-19 by income, race/ethnicity, and household crowding: US county vs ZIP code analyses. J Public Health Management Policy (in press).

[32] Felsen M. OSHA is failing to protect workers. The Shelbyville News 2020 August 13
(https://www.shelbynews.com/osha-is-failing-to-protect-workers/article_391496df-7856-5f8982da-2d0a6afd36d0.html).

[33] Campbell AF, Yerardi J. Amid the COVID-19 pandemic, the U.S. Occupational Safety and Health Administration has been cutting back even more. The Center for Public Integrity and Vox 2020 August 18 (https://publicintegrity.org/politics/system-failure/deaths-cutbacks-workplacesafety-inspections-osha/).

[34] The Office of (Michigan) Governor Gretchen Whitmer. Executive Order 2020-36 (COVID-19)— Protecting workers who stay home, stay safe when they or their close contacts are sick. 2020 (https://www.michigan.gov/whitmer/0,9309,7-387-90499_90705-524136--,00.html).

[35] Virginia Department of Labor and Industry. Press release—Virginia Occupational Safety and Health program publishes emergency temporary standard for infectious disease prevention: SARS-CoV-2 virus that causes covid-19. 2020 July 27 (https://www.doli.virginia.gov/wp-

content/uploads/2020/07/COVID-19-ETS-News-Release-DRAFT-7.23.2020-1.pdf).

[36] State of California Dept of Industrial Relations. New release— Cal/OSHA issues citations to multiple employers for COVID-19 violations. Release number 2020-76; 2020 September 4 (https://www.dir.ca.gov/DIRNews/2020/2020-76.html).

[37] Krieger N, Waterman PD, Chen JT. COVID-19 and overall mortality inequities in the surge in death rates by ZIP Code characteristics: Massachusetts, January 1 to May 19, 2020. Am J Public Health (in press).

[38] Leon DA, Shkolnikov VM, Smeeth L, et al. COVID-19: a need for real-time monitoring of weekly excess deaths (letter). Lancet 2020 May 2;395(10234):e81 (https://doi.org/10.1016/S0140-6736(20)30933-8) (Epub ahead of print).

[39] Preliminary Estimate of Excess Mortality During the COVID-19 Outbreak—New York City,

March 11–May 2, 2020. Morb Mortal Wkly Rep 2020;69:603-605

(http://dx.doi.org/10.15585/mmwr.mm6919e5).

[40] CDC. Weekly counts of deaths by jurisdiction and age group

(https://data.cdc.gov/NCHS/Weekly-counts-of-deaths-by-jurisdiction-and-age-gr/y5bj-9g5w/).

[41] CDC. CDC Wonder. Bridged-race population estimates 1990-2019

(https://wonder.cdc.gov/Bridged-Race-v2019.html).

[42] Rothman KJ, Greenland S. Modern epidemiology, 2nd ed. Chapter 14. Philadelphia, PA: Lippincott-Raven, 1988.

[43] CDC National Center for Health Statistics. Excess deaths associated with COVID-19: provisional death counts for coronavirus disease (COVID-19)

(https://www.cdc.gov/nchs/nvss/vsrr/covid19/excess_deaths.htm).

[44] Rucker P, Wan W. Trump projects up to 240,000 coronavirus deaths in the U.S., even with mitigation efforts. Washington Post 2020 April 1

(https://www.washingtonpost.com/politics/trump-white-house-projects-up-to-240000coronavirus-deaths-in-us-even-with-mitigation-efforts/2020/03/31/62df5344-7367-11ea-87da-77a8136c1a6d story.html).

[45] Slisco A. Trump advisers doubt White House's coronovirus deaths estimate, expert says it appears rushed: report. Newsweek 2020 April 3 (https://www.newsweek.com/trump-advisers-doubt-white-houses-coronavirus-deaths-estimate-expert-says-it-appeared-rushed-1495908).

[46] Osborne H. Why the COVID-19 death forecasts are wrong. Newsweek 2020 April 21

(https://www.newsweek.com/why-covid-19-death-forecasts-are-wrong-1499243).

[47] Xu JQ, Murphy SL, Kochanek KD, et al. Mortality in the United States, 2018. NCHS Data Brief,

no 355. Hyattsville, MD: National Center for Health Statistics. 2020

(https://www.cdc.gov/nchs/products/databriefs/db355.htm).

[48] Australian Bureau of Statistics. Provisional Mortality Statistics, Jan–May 2020

(https://www.abs.gov.au/ausstats/abs@.nsf/mf/3303.0.55.004).

[49] Merkler AE, Parikh NS, Mir S, et al. Risk of ischemic stroke in patients with coronavirus disease 2019 (COVID-19) vs patients with influenza. JAMA Neuro 2020 July 2

(doi:10.1001/jamaneurol.2020.2730).

[50] Yonker LM, Neilan AM, Bartsch Y, et al. Pediatric SARS-CoV-2: clinical presentation, infectivity,

and immune responses. J Pediatr 2020 Aug 18;S0022-3476(20)31023-4 (doi:

10.1016/j.jpeds.2020.08.037) (Epub ahead of print).

[51] US Census. Quick Facts: Massachusetts. Population estimates, July 1, 2019

(https://www.census.gov/quickfacts/fact/table/MA/PST045219).

[52] Massachusetts Department of Public Health. Cancer incidence and mortality in Massachusetts 2012-2016: statewide report. 2020 August (https://www.mass.gov/doc/cancer-incidence-and-mortality-in-massachusetts-2012-2016-with-a-special-overview-mass-in-0/download).

[53] US National Cancer Institute. Cancer stat facts: common cancer sites. New cancer cases, 2020

(https://seer.cancer.gov/statfacts/html/common.html).

[54] Godlee F. Living with covid-19. BMJ 2020;370:m3392 (doi:

https://doi.org/10.1136/bmj.m3392).

[55] Oppel Jr RA, Gebeloff R, Lai KKR, et al. The fullest look yet at the racial inequity of coronavirus. New York Times 2020 July 5 (https://www.nytimes.com/interactive/2020/07/05/us/coronaviruslatinos-african-americans-cdc-data.html).

[56] Bassett MT, Chen JT, Krieger N. The unequal toll of COVID-19 mortality by age in the United States: quantifying racial/ethnic disparities. Harvard Center for Population and Development Studies Working Paper Series 2020;19(3) (https://cdn1.sph.harvard.edu/wpcontent/uploads/sites/1266/2020/06/20_Bassett-Chen-Krieger_COVID-19_plus_age_workingpaper_0612_Vol-19_No-3_with-cover.pdf). [57] CDC. Deaths involving coronavirus disease 2019 (COVID-19) by race and Hispanic origin group and age, by state (https://data.cdc.gov/NCHS/Deaths-involving-coronavirus-disease-2019-COVID-19/ks3g-spdg).

[58] Servick K. 'Huge hole' in COVID-19 testing data makes it harder to study racial disparities.Science/AAAS Science Magazine News 2020 July 10

(https://www.sciencemag.org/news/2020/07/huge-hole-covid-19-testing-data-makes-it-harder-study-racial-disparities).

[59] Lieberman-Cribbin W, Tuminello S, Flores RM, et al. Disparities in COVID-19 testing and positivity in New York City. Am J Prev Med 2020:S0749-3797(20)30263-4

(doi:10.1016/j.amepre.2020.06.005) (Epub ahead of print).

[60] Chen Y, Freedman ND, Rodriquez EJ, et al. Trends in premature deaths among adults in the

United States and Latin America. JAMA Netw Open 2020;3(2):e1921085

(doi:10.1001/jamanetworkopen.2019.21085).

[61] Cunningham TJ, Croft B, Liu Y, et al. Vital Signs: racial disparities in age-specific mortality

among Blacks or African-Americans – United States, 1999-2015. Morb Mortal Wkly Rep 2017 May

2 (https://www.cdc.gov/mmwr/volumes/66/wr/mm6617e1.htm).

[62] Worobey M, Pekar J, Larsen BB, et al. The emergence of SARS-CoV-2 in Europe and North

America. Science/AAAS Science Magazine News 2020 September 10

(DOI:10.1126/science.abc8169).

[63] Peak CM, Kahn R, Grad YH, et al. Individual quarantine versus active monitoring of contacts for the mitigation of COVID-19: a modelling study. The Lancet Infec Dis 2020;20(9):1025-1033 (https://www.thelancet.com/journals/laninf/article/PIIS1473-3099(20)30361-3/fulltext).

[64] Michaels D, Wagner GR. Occupational Safety and Health Administration (OSHA) and worker safety during the COVID-19 pandemic. JAMA 2020 September 16 (doi:10.1001/jama.2020.16343).

Table 1. Estimation of excess deaths during the US COVID-19 pandemic, comparing weekly 2020 death counts to the corresponding average annual 2015-2019 deaths, using 4 different methods^a (source: Centers for Disease Control and Prevention)

	Crude	(050) (0)		Age- standardized excess rate per 100,000 person-	(050) (0)		Age- standardized cumulative incidence proportion per 100,000	(055)(01)		
Method	excess	(95% CI)		years ^₅	(95% CI)		population	(95% CI)		
1	263,375	(262,369	, 264,381)	109.0	(108.2	, 109.8)	66.8	(66.3	, 67.3)	
2	272,449	(271,426	, 273,472)	112.7	(111.9	, 113.5)	69.1	(68.6	, 69.6)	
3	231,049	(230,107	, 231,991)	95.1	(94.4	, 95.9)	58.3	(57.9	, 58.8)	
4	246,758	(245,784	, 247,731)	101.8	(101.0	, 102.5)	62.4	(61.9	, 62.9)	

^a Method 1: sum up over weeks and then age-standardize using the direct method

Method 2: count only age strata and weeks where the excess is greater than zero, and set weeks where the excess in the age stratum is less than zero to zero

Method 3: instead of comparing to the average deaths in 2015-2019, we compare to the 95% upper bound on the average deaths for 2015-2019

Method 4: we compare to the upper bound on the average deaths for 2015-2019, but once again ignore weeks where the excess in the age stratum is less than zero

^b Age-standardized excess rate per 100,000 person-years is computed based on dividing the age-specific excess count of deaths under the four methods by the age-specific population person-time (taking into account the age-specific 2020 population counts and the elapsed time since January 1, 2020), weighting by the year 2000 standard million, and summing over age categories. ^c Age-standardized cumulative incidence proportion per 100,000 population (i.e. a risk per capita) is computed based on dividing the age-specific excess count of deaths under the four methods by the age-specific population count in 2020, weighting by the year 2000 standard million, and summing over age categories.

Figure 1 National OSHA complaints and COVID—19 deaths per million (7 Day Average)





